

**TOPIC 9**  
**VECTOR ALGEBRA**  
**SCHEMATIC DIAGRAM**

Topic	Concept	Degree of importance	Reference
			NCERT Text Book Edition 2007
Vector algebra	(i) Vector and scalars	*	Q2 pg428
	(ii) Direction ratio and direction cosines	*	Q 12,13 pg 440
	(iii) Unit vector	* *	Ex 6,8 Pg 436
	(iv) Position vector of a point and collinear vectors	* *	Q 15 Pg 440 , Q 11Pg440 , Q 16 Pg448
	(v) Dot product of two vectors	**	Q6 ,13 Pg445
	(vi) Projection of a vector	* * *	Ex 16 Pg 445
	(vii) Cross product of two vectors	* *	Q 12 Pg458
	(viii) Area of a triangle	*	Q 9 Pg 454
	(ix) Area of a parallelogram	*	Q 10 Pg 455

**SOME IMPORTANT RESULTS/CONCEPTS**

\* Position vector of point A(x, y, z) =  $\vec{OA} = x\hat{i} + y\hat{j} + z\hat{k}$

\* If A(x<sub>1</sub>, y<sub>1</sub>, z<sub>1</sub>) and point B(x<sub>2</sub>, y<sub>2</sub>, z<sub>2</sub>) then  $\vec{AB} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$

\* If  $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$  ;  $|\vec{a}| = \sqrt{x^2 + y^2 + z^2}$

\* Unit vector parallel to  $\vec{a} = \frac{\vec{a}}{|\vec{a}|}$

\* Scalar Product (dot product) between two vectors :  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$  ;  $\theta$  is angle between the vectors

\*  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

\* If  $\vec{a} = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$  and  $\vec{b} = a_2\hat{i} + b_2\hat{j} + c_2\hat{k}$  then  $\vec{a} \cdot \vec{b} = a_1a_2 + b_1b_2 + c_1c_2$

\* If  $\vec{a}$  is perpendicular to  $\vec{b}$  then  $\vec{a} \cdot \vec{b} = 0$

\*  $\vec{a} \cdot \vec{a} = |\vec{a}|^2$

\* Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

\* Vector product between two vectors :

$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}$  ;  $\hat{n}$  is the normal unit vector which is perpendicular to both  $\vec{a}$  &  $\vec{b}$

\*  $\hat{n} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$

\* If  $\vec{a}$  is parallel to  $\vec{b}$  then  $\vec{a} \times \vec{b} = 0$

\* Area of triangle (whose sides are given by  $\vec{a}$  and  $\vec{b}$ ) =  $\frac{1}{2} |\vec{a} \times \vec{b}|$

\* Area of parallelogram (whose adjacent sides are given by  $\vec{a}$  and  $\vec{b}$ ) =  $|\vec{a} \times \vec{b}|$

\* Area of parallelogram (whose diagonals are given by  $\vec{a}$  and  $\vec{b}$ ) =  $\frac{1}{2} |\vec{a} \times \vec{b}|$

## ASSIGNMENTS

*(i) Vector and scalars, Direction ratio and direction cosines & Unit vector*

### LEVEL I

1. If  $\vec{a} = \hat{i} + \hat{j} - 5\hat{k}$  and  $\vec{b} = \hat{i} - 4\hat{j} + 3\hat{k}$  find a unit vector parallel to  $\vec{a} + \vec{b}$
2. Write a vector of magnitude 15 units in the direction of vector  $\hat{i} - 2\hat{j} + 2\hat{k}$
3. If  $\vec{a} = \hat{i} + \hat{j} - \hat{k}$  ;  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  ;  $\vec{c} = -\hat{i} + \hat{j} + \hat{k}$  find a unit vector in the direction of  $\vec{a} + \vec{b} + \vec{c}$
4. Find a unit vector in the direction of the vector  $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$  [ CBSE 2011]
5. Find a vector in the direction of vector  $\vec{a} = \hat{i} - 2\hat{j}$ , whose magnitude is 7

### LEVEL II

1. Find a vector of magnitude 5 units, perpendicular to each of the vectors  $(\vec{a} + \vec{b})$ ,  $(\vec{a} - \vec{b})$  where

$$\vec{a} = \hat{i} + \hat{j} + \hat{k} \text{ and } \vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}.$$

- If the sum of two unit vectors is a unit vector, show that the magnitude of their difference is  $\sqrt{3}$ .
- If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$  and  $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$ , find a vector of magnitude 6 units which is parallel to the vector  $2\vec{a} - \vec{b} + 3\vec{c}$

### LEVEL – III

- If a line make  $\alpha, \beta, \gamma$  with the X - axis , Y- axis and Z – axis respectively, then find the value of  $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma$
- For what value of p, is  $(\hat{i} + \hat{j} + \hat{k}) p$  a unit vector?
- What is the cosine of the angle which the vector  $\sqrt{2} \hat{i} + \hat{j} + \hat{k}$  makes with Y-axis
- Write the value of p for which  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$  are parallel vectors.

### (ii) Position vector of a point and collinear vectors

#### LEVEL – I

- Find the position vector of the midpoint of the line segment joining the points  $A(5\hat{i} + 3\hat{j})$  and  $B(3\hat{i} - \hat{j})$ .
- In a triangle ABC, the sides AB and BC are represents by vectors  $2\hat{i} - \hat{j} + 2\hat{k}$ ,  $\hat{i} + 3\hat{j} + 5\hat{k}$  respectively. Find the vector representing CA.
- Show that the points (1,0), (6,0), (0,0) are collinear.

#### LEVEL – II

- Write the position vector of a point R which divides the line joining the points P and Q whose position vectors are  $\hat{i} + 2\hat{j} - \hat{k}$  and  $-\hat{i} + \hat{j} + \hat{k}$  respectively in the ratio 2 : 1 externally.
- Find the position vector of a point R which divides the line joining two points P and Q whose position vectors are  $(2\vec{a} + \vec{b})$  and  $(\vec{a} - 3\vec{b})$  respectively, externally in the ratio 1:2. Also, show that P is the mid-point of the line segment RQ

### (iii) Dot product of two vectors

#### LEVEL – I

- Find  $\vec{a} \cdot \vec{b}$  if  $\vec{a} = 3\hat{i} - \hat{j} + 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} + 3\hat{k}$ .

2. If  $|\vec{a}| = \sqrt{3}$ ,  $|\vec{b}| = 2$  and  $\vec{a} \cdot \vec{b} = \sqrt{6}$ . Then find the angle between  $\vec{a}$  and  $\vec{b}$ .
3. Write the angle between two vectors  $\vec{a}$  and  $\vec{b}$  with magnitudes  $\sqrt{3}$  and 2 respectively having  $\vec{a} \cdot \vec{b} = \sqrt{6}$  [CBSE 2011]

### LEVEL – II

1. The dot products of a vector with the vectors  $\hat{i} - 3\hat{j}$ ,  $\hat{i} - 2\hat{j}$  and  $\hat{i} + \hat{j} + 4\hat{k}$  are 0, 5 and 8 respectively. Find the vectors.
2. If  $\vec{a}$  and  $\vec{b}$  are two vectors such that  $|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$ , then what is the angle between  $\vec{a}$  and  $\vec{b}$ .
3. If  $\vec{a} = 2\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$  and  $\vec{c} = 3\hat{i} + \hat{j}$  are such that  $\vec{a} + \lambda\vec{b}$  is perpendicular to  $\vec{c}$ , find the value of  $\lambda$ .

### LEVEL – III

1. If  $\vec{a}$  &  $\vec{b}$  are unit vectors inclined at an angle  $\theta$ , prove that  $\sin \frac{\theta}{2} = \frac{1}{2} |\vec{a} - \vec{b}|$ .
2. If  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ , then find the angle between  $\vec{a}$  and  $\vec{b}$ .
3. For what values of  $\lambda$ , vectors  $\vec{a} = 3\hat{i} - 2\hat{j} + 4\hat{k}$  and  $\vec{a} = \lambda\hat{i} - 4\hat{j} + 8\hat{k}$  are  
(i) Orthogonal (ii) Parallel
4. Find  $|\vec{x}|$ , if for a unit vector  $\vec{a}$ ,  $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 15$ .
5. If  $\vec{a} = 5\hat{i} - \hat{j} + 7\hat{k}$  and  $\vec{b} = \hat{i} - \hat{j} + \mu\hat{k}$ , find  $\mu$ , such that  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  are orthogonal.
6. Show that the vector  $2\hat{i} - \hat{j} + \hat{k}$ ,  $-3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  form sides of a right angled triangle.
7. Let  $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$ ,  $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$ . Find a vector  $\vec{d}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and  $\vec{c} \cdot \vec{d} = 18$ .
8. If  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are three mutually perpendicular vectors of equal magnitudes, prove that  $\vec{a} + \vec{b} + \vec{c}$  is equally inclined with the vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$ .
9. Let  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  be three vectors such that  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$ ,  $|\vec{c}| = 5$  and each of them being perpendicular

to the sum of the other two, find  $\left| \vec{a} + \vec{b} + \vec{c} \right|$ .

(iv) *Projection of a vector*

**LEVEL – I**

1. Find the projection of  $\vec{a}$  on  $\vec{b}$  if  $\vec{a} \cdot \vec{b} = 8$  and  $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$ .
2. Write the projection of the vector  $\hat{i} - \hat{j}$  on the vector  $\hat{i} + \hat{j}$  [CBSE 2011]
3. Find the angle between the vectors  $\hat{i} - 2\hat{j} + 3\hat{k}$  and  $3\hat{i} - 2\hat{j} + \hat{k}$
4. Find the projection of the vector  $\hat{i} + 3\hat{j} + 7\hat{k}$  on the vector  $7\hat{i} - \hat{j} + 8\hat{k}$

**LEVEL – II**

1. Three vertices of a triangle are A(0, -1, -2), B(3, 1, 4) and C(5, 7, 1). Show that it is a right angled triangle. Also find the other two angles.
2. Show that the angle between any two diagonals of a cube is  $\cos^{-1}\left(\frac{1}{3}\right)$ .

3. If  $\vec{a}, \vec{b}, \vec{c}$  are non-zero and non-coplanar vectors, prove that  $\vec{a} - 2\vec{b} + 3\vec{c}, -3\vec{b} + 5\vec{c}$  and  $-2\vec{a} + 3\vec{b} - 4\vec{c}$  are also coplanar

**LEVEL – III**

1. If a unit vector  $\vec{a}$  makes angles  $\pi/4$  with  $\hat{i}$ ,  $\pi/3$  with  $\hat{j}$  and an acute angle  $\theta$  with  $\hat{k}$ , then find the component of  $\vec{a}$  and angle  $\theta$ .
2. If  $\vec{a}, \vec{b}, \vec{c}$  are three mutually perpendicular vectors of equal magnitudes, prove that  $\vec{a} + \vec{b} + \vec{c}$  is equally inclined with the vectors  $\vec{a}, \vec{b}, \vec{c}$ .
3. If with reference to the right handed system of mutually perpendicular unit vectors  $\hat{i}, \hat{j}$ , and  $\hat{k}$ ,  $\vec{\alpha} = 3\hat{i} - \hat{j}$ ,  $\vec{\beta} = 2\hat{i} + \hat{j} - 3\hat{k}$  then express  $\vec{\beta}$  in the form of  $\vec{\beta}_1 + \vec{\beta}_2$ , where  $\vec{\beta}_1$  is parallel to  $\vec{\alpha}$  and  $\vec{\beta}_2$  is perpendicular to  $\vec{\alpha}$ .
4. Show that the points A, B, C with position vectors  $\vec{a} = 3\hat{i} - 4\hat{j} - 4\hat{k}$ ,  $\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} - 3\hat{j} - 5\hat{k}$  respectively form the vertices of a right angled triangle.

5. If  $\vec{a}$  &  $\vec{b}$  are unit vectors inclined at an angle  $\theta$ , prove that

$$(i) \sin \frac{\theta}{2} = \frac{1}{2} |\vec{a} - \vec{b}| \quad (ii) \tan \frac{\theta}{2} = \frac{|\vec{a} - \vec{b}|}{|\vec{a} + \vec{b}|}$$

(vii) *Cross product of two vectors*

### LEVEL – I

1. If  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$  and  $\vec{a} \cdot \vec{b} = 9$ . Find  $|\vec{a} \times \vec{b}|$

2. Find  $|\vec{a} \times \vec{b}|$ , if  $\vec{a} = \hat{i} - 7\hat{j} + 7\hat{k}$  and  $\vec{b} = 3\hat{i} + 2\hat{j} + 2\hat{k}$

3. Find  $|\vec{x}|$ , if  $\vec{p}$  is a unit vector and  $(\vec{x} - \vec{p}) \cdot (\vec{x} + \vec{p}) = 80$ .

4. Find  $p$ , if  $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + 3\hat{j} + p\hat{k}) = \vec{0}$ .

### LEVEL – II

1. Find  $\lambda$ , if  $(2\hat{i} + 6\hat{j} + 14\hat{k}) \times (\hat{i} - \lambda\hat{j} + 7\hat{k}) = \vec{0}$ .

2. Show that  $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2(\vec{a} \times \vec{b})$

3. Find the angle between two vectors  $\vec{a}$  and  $\vec{b}$  if  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$  and  $|\vec{a} \times \vec{b}| = 6$ .

4. Let  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  be unit vectors such that  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} = 0$  and the angle between  $\vec{b}$  and  $\vec{c}$  is  $\pi/6$ , prove that  $\vec{a} = \pm 2(\vec{a} \times \vec{b})$ .

### LEVEL – III

1. Find the value of the following:  $\hat{i} \cdot (\hat{j} \times \hat{k}) + \hat{i} \cdot (\hat{i} \times \hat{k}) + \hat{k} \cdot (\hat{i} \times \hat{j})$

2. Vectors  $\vec{a}$  and  $\vec{b}$  are such that  $|\vec{a}| = \sqrt{3}$ ,  $|\vec{b}| = \frac{2}{3}$ , and  $\vec{a} \times \vec{b}$  is a unit vector. Write the

angle between  $\vec{a}$  and  $\vec{b}$

3. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{b} = \hat{j} - \hat{k}$ , find a vector  $\vec{c}$  such that  $\vec{a} \times \vec{c} = \vec{b}$  and

$$\vec{a} \cdot \vec{c} = 3.$$

4. If  $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$  and  $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$  show that  $(\vec{a} - \vec{d})$  is parallel to  $\vec{b} - \vec{c}$ , where

$$\vec{a} \neq \vec{d} \text{ and } \vec{b} \neq \vec{c}.$$

5. Express  $2\hat{i} - \hat{j} + 3\hat{k}$  as the sum of a vector parallel and perpendicular to  $2\hat{i} + 4\hat{j} - 2\hat{k}$ .

(viii) *Area of a triangle & Area of a parallelogram*

#### LEVEL – I

1. Find the area of Parallelogram whose adjacent sides are represented by the vectors

$$\vec{a} = 3\hat{i} + \hat{j} - 2\hat{k} \text{ and } \vec{b} = \hat{i} - 3\hat{j} + 4\hat{k}.$$

2. If  $\vec{a}$  and  $\vec{b}$  represent the two adjacent sides of a Parallelogram, then write the area of parallelogram in terms of  $\vec{a}$  and  $\vec{b}$ .

3. Find the area of triangle having the points A(1,1,1), B(1,2,3) and C(2,3,1) as its vertices.

#### LEVEL – II

1. Show that the area of the Parallelogram having diagonals  $(3\hat{i} + \hat{j} - 2\hat{k})$  and  $(\hat{i} - 3\hat{j} + 4\hat{k})$  is  $5\sqrt{3}$  Sq units.

2. If  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are the position vectors of the vertices of a  $\Delta ABC$ , show that the area of the  $\Delta ABC$  is

$$\frac{1}{2} \left| \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} \right|.$$

3. Using Vectors, find the area of the triangle with vertices A(1,1,2), B(2,3,5) and C(1,5,5)  
[CBSE 2011]

#### Questions for self evaluation

1. The scalar product of the vector  $\hat{i} + \hat{j} + \hat{k}$  with the unit vector along the sum of vectors

$$2\hat{i} + 4\hat{j} - 5\hat{k} \text{ and } \lambda\hat{i} + 2\hat{j} + 3\hat{k} \text{ is equal to one. Find the value of } \lambda.$$

2. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  be three vectors such that  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$ ,  $|\vec{c}| = 5$  and each one of them being perpendicular to the sum of the other two, find  $|\vec{a} + \vec{b} + \vec{c}|$ .

3. If  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ , then find the angle between  $\vec{a}$  and  $\vec{b}$ .

4. Dot product of a vector with  $\hat{i} + \hat{j} - 3\hat{k}$ ,  $\hat{i} + 3\hat{j} - 2\hat{k}$ , and  $2\hat{i} + \hat{j} + 4\hat{k}$  are 0, 5, 8 respectively.  
Find the vector.

5. Find the components of a vector which is perpendicular to the vectors  $\hat{i} + 2\hat{j} - \hat{k}$  and  $3\hat{i} - \hat{j} + 2\hat{k}$ .