# Downloaded from www.studiestoday.com

#### **Chapter 10**

#### STRAIGHT LINES

**SLOPE OF A LINE** :  $m = tan\theta$  if  $\theta$  is the angle of inclination.

m =  $\frac{y_2 - y_1}{x_2 - x_1}$  if  $(x_1, y_1)$  and  $(x_2, y_2)$  are two points on the line.

SLOPE of a horizontal line is 0 and vertical line is not defined.

If  $m_1$  and  $m_2$  are slopes of  $L_1$  and  $L_2$  respectively.

 $L_1 \quad \parallel \quad L_2 \rightarrow \quad m_1 = m_2$ 

 $L_1 \perp L_2 \quad {\scriptstyle \rightarrow} \quad m_1 \; x \; m_2 \; = \; -1$ 

Acute angle between  $L_1$  and  $L_2$ 

 $\tan \theta = \left| \frac{m^2 - m^1}{1 + m^1 \times m^2} \right| as \ 1 + m_1 m_2 \neq 0 \text{ and } \text{ the obtuse angle}$  $\emptyset = 180 - \theta.$ 

#### EQUATION OF STRAIGHT LINE

x-axis y-axis

# y = 0x = 0

- $|| \text{ to } x\text{-axis} \qquad \rightarrow \qquad y = t$
- $\parallel \text{ to y-axis } \longrightarrow x = a$

Having slope m and making an intercept c on y-axis $\rightarrow$  y = mx+c

Making intercepts a and b on the x-axis and y-axis  $\rightarrow \frac{x}{a} + \frac{y}{b} = 1$ 

passing through  $(x_1, y_1)$  and  $(x_2, y_2) \rightarrow y - y_1 = \frac{y_2 - y_1}{x_2 - x_1} (x - x_1)$ 

Having normal distance from orgin P and angle between the normal and positive x-axis  $\omega \rightarrow x \cos \omega + y \sin \omega = P$ .

General form  $\rightarrow$  Ax + By + C = 0

Distance of a point  $(x_1, y_1)$  from a line ax + by + c = 0 is  $\left|\frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}\right|$ 

### Downloaded from www.studiestoday.com

## Downloaded from www.studiestoday.com

#### **TEXT BOOK QUESTIONS**

$\rightarrow$	Exercise	10.1	$\rightarrow$	Qns 5,8,9
---------------	----------	------	---------------	-----------

- \*  $\rightarrow$  Exercise 10.2  $\rightarrow$  Qns 7,8,9,10,11,16
- \*  $\rightarrow$  Exercise 10.3  $\rightarrow$  Qns 3,4,5,7,8,9,10,12,16
- \* → MiscExercis→ Qns 1,6,7,8,9,12,14,15,23
- \*\*  $\rightarrow$  Exercise 10.1  $\rightarrow$  Qns 11,13
- \*\*  $\rightarrow$  Exercise 10.2  $\rightarrow$  Qns 12,13,15,18,20
- \*\*  $\rightarrow$  Exercise 10.3  $\rightarrow$  Qns 13,14,17,18
- \*\*  $\rightarrow$  Misc Exercise  $\rightarrow$  Qns 3,4,11,18,19
- \*\*  $\rightarrow$  Example  $\rightarrow$  2,3,13,14,15,17,19,20,23

Misc Example  $\rightarrow 23$ 

\*

#### **EXTRA/ HOT QUESTIONS**

1. Find the equation of the line through (4,-5) and parallel to the line joining the points (3,7) & (-2,4).

(Ans.3x-5y-37=0)

- 2. If A(1,4), B(2,-3) and C(-1,-2) are the vertices of a triangle ABC . find
  - a) The equation of the median through A
  - b) The equation of the altitude through A
  - c) The right bisector of side BC
- 3. Find the equation of the straight line which passes through (3,-2) and cuts off positive intercepts on the x axis and y axis which are in the ratio 4:3
- 4. Reduce the equation 3x-2y+4 = 0 to intercept form. Hence find the length of the segment intercepted between the axes.
- 5. Find the image of the point (1,2) in the line x-3y+4=0
- 6. If the image of the point (2,1) in a line is (4,3) .Find the equation of the line.
- 7. Find the equation of a line passing through the point (-3,7) and the point of intersection of the lines 2x-3y+5 = 0 and 4x+9y = 7.

(Ans.8x+3y+3=0)

### Downloaded from www.studiestoday.com

# Downloaded from www.studiestoday.com

8. Find the equation of straight lines which are perpendicular to the line

12x+5y = 17 and at a distance of 2 units from the point (-4,1)

(ans. 5x-12y+6=0 & 5x-12y+58=0)

9. The points A(2,3) B(4,-1) & C(-1,2) are the vertices of a triangle. Find the length of perpendicular from A to BC and hence the area of ABC (Ans.  $\frac{14}{\sqrt{34}}$  units & 7 sq.units)

10. Find the equation of straight line whose intercepts on the axes are thrice as long as those made by 2x + 11y = 6

Downloaded from www.studiestoday.com

(Ans.2x+11y=18)