


Let us remember :

Draw bar graph from the given information :

Name of fruit	Chikoo	Grapes	Watermelon	Mango	Apple
Sale (kg)	18	36	45	54	27


```
Title : .....
```



Downloaded from https:// www.studiestoday.com

1 : Pie Graph

- Let us learn new :
 - If the information given in the bar-graph is shown in a single circle, it is called a pie-graph.
 - To show the information in a circle, the information should be transformed into degree measure.
 - The measure of all the angles at the center of the circle is 360°. See in the figure here.
 - When the information other than measure of angles is to be shown, the base of 360° is taken.



• Let's construct the pie-graph with the help of the information given in the bar-graph.

Sr. No.	Fruit	kg
(1)	Chikoo	18
(2)	Grapes	36
(3)	Watermelon	45
(4)	Mango	54
(5)	Apple	27

- Total sale of fruits (in kg) = 180 kg
- The complete circle is shown with 360°.

Total sale 180 kg : 360°

∴ 18 kg : (?)


$$=\frac{18\times360}{180}=36^{\circ}$$

- Transfer of information into percentage :
 - Sale of Chikoo out of total 180 kg = 18 kg
- :. Sale of Chikoo out of total 100 kg = (?)

$$=\frac{18 \times 100}{180} = 10 \text{ kg}$$

Sale of Chikoo was 10 %.





- You have seen that for finding degree measure, base of 360° is taken and for finding percentage, base of 100 is taken.
- Fill the given Table by calculating the information in the above manner. Later, with the help of your teacher, prepare a pie-graph in a computer.

Sr. No.	Fruit	kg	Degree	Percentage
(1)	Chikoo	18	36 ⁰	10 %
(2)	Grapes	36		
(3)	Watermelon	45		
(4)	Mango	54		
(5)	Apple	27		

The pictorial representation of the information in a circular area is known as pie-graph.

- Where have you find such pie-graphs ? What kind of different information is shown with the pie-graph ?
- Discuss this with your teacher and note down about the uses of pie-graph in your note-book.




1. Some questions are given based on the below given pie graph. Study the pie graph and answer the questions with calculation in your notebook :

The contribution given by the students of Std. 7 for the celebration of Teacher's day is shown in the pie graph given below :

- (1) In all how many rupees were collected as fund ?
- (2) How much contribution was given by Jafar ?
- (3) Calculate the degree measure of the contribution by Anuj and measure it in the figure.
- (4) Who contributed the least fund ?


MATHEMATICS



STD.7



- 2. As shown in the pie graph, Ajaybhai bought 3600 kilogram grains from the market for his shop during the day, answer the following questions :
 - (1) How much percentage is cotton more than wheat ?
 - (2) How many kilograms of sesame was bought ?
 - (3) Calculate the degree measure of the part of cotton and measure it.
 - (4) How many kilogram of Castor-seeds are there ?



- 3. Based on the pie graph showing 50 runs scored in the first over in a cricket match, answer the following questions :
 - (1) In which over maximum runs were scored ?
 - (2) In which over, not a single run was scored ?
 - (3) How many runs were scored in the third over ?
 - (4) In which over, 28 percentage of runs were scored ?






1. The monthly income of Anjaliben is ₹ 7200. Study the pie-graph of the monthly budget prepared by her and answer the following questions :

- (1) How many percentage of expenditure is done for education ?
- (2) What is the budget for savings in terms of money ?
- (3) How many degree measure is shown by the expenditure on clothes ?
- (4) On what is the maximum expenditure done ?

MATHEMATICS



STD.7




2 Integers

Let us remember :


Students, we have learnt about integers and its addition and subtraction. Let us remember them again.

1. Fill in the blanks with the help of number line :



- (1) The cartoon is standing on digit.
- (2) integers are there on the left side of zero.
- (3) Positive integers are on the side of zero (0).
- (4) There is integer, 4 units right to zero.
- (5) There is integer, 4 units left to zero.

2. Fill in the blanks with the help of number line :







Thus, the numerical value of a number obtained by disregarding the sign prefixed to the number is called its absolute value.

To denote the absolute value the sign '| ' (modulus) is used.

That is, the absolute value of 5 = 5

It is written symbolically |5| = 5 (Read as : modulus five is equal to five) The absolute value of (-6) = 6

It is written symbolically |-6| = 6 (Read as : modulus minus six is equal to six) The absolute value of 0 = 0

It is written symbolically |0| = 0 (Read as : modulus zero is equal to zero) The absolute value of no number can be negative.

Zero, positive numbers and negative numbers are included in integers.
 Properties of addition of integers :

We have learnt about properties of addition of whole number in fifth standard. Now, in the same way, we are going to learn about properties of addition of integers.

1. Look, understand and complete :

(-5)	+	4	=	(-1)	
integer		integer		integer	
(-7)	+	(-8)	=	(—15)	
integer		integer		integer	
	+	6	=	2	
integer		integer		integer	
(-7)	+	(-1)	=		
integer		integer		integer	

It is clear from the above Table that the addition of any two integers is also an integer.




Downloaded from https:// www.studiestoday.com

2 : Integers

• Activity 1 :

- Prepare small _____ flash cards from a thick paper.
- Write 0 to 9 and (-1) to (-9) on the cards.
- Prepare flash cards with signs of +, and =.
- Gather all the cards in one box.
- Now, from that take any two digit cards and put + or sign card between them.
- Then arrange the answer card after the card with = sign.
- Example :



- If the position of the digit card is changed, what will be the answer ?
- Do this activity, three to four times by changing the flash cards.
- What difference do you find in the answer ? Note down with practice.
- If the same activity is done using subtraction, what can be seen ?
- Is the answer same or different ?
- 2. Look, understand and complete :

(-4) + 5 = 5 + (-4) = 1 3 + (-7) = (-7) + 3 = (-4) $(-8) + \dots = (-6) + (-8) = (-14)$ $(-10) + 5 = 5 + \dots = (-5)$

It is clear from the above examples that if we add any integers in any order the result is same.



		2 : Integers	
3.	Look, understand and	complete :	
	[(-3) + 6] + 7 or	(-3) + [6 + 7] or	[(-3) + 7] + 6
	= 3 + 7	= (-3) + (13)	= 4 + 6
	= 10	= 10	= 10
	Complete as above :		
	[4 + (-5)] + 7 or	4 + [(-5) + 7] or	[4 + 7] + (-5)
	= (-1) +	= 4 +	= + (-5)
	= 6	= 6	= 6
	Thus, to do the ad	dition of any three into	egers, taking a group of first

Thus, to do the addition of any three integers, taking a group of first two integers and then adding the third integer to the sum, the result will be the same.

4. Look, understand and complete :

$$7 + 0 = 7$$

(-8) + 0 =
0 + 9 = 9
0 + = (-15)

If we add zero to any integer, then we get the same integer.

Thus, 0 is called the neutral number (identity) for addition.

- 5. Look, understand and complete :
 - 7 + (-7) = 0(-8) + 8 = 15 + = 0 (-12) + = 0

Thus, if addition of any two integers is zero, then each one is opposite integer (or additive inverse) of each other.

That is to say opposite number of 7 = (-7). The opposite number of (-9) = 9and 0 + 0 = 0, that is to say opposite number of 0 = 0.



2 : Integers

Addition and subtraction of integers :

We know that if the number added decreases by 1 the result also decreases by 1.

Addition of positive integer or zero to positive integer gives positive integer.	3 + 3 = 6 3 + 2 = 5 3 + 1 = 4 3 + 0 = 3	(-3) + 6 = 3 $(-3) + 5 = 2$ $(-3) + 4 = 1$ $(-3) + 3 = 0$ $(-3) + 2 = (-1)$	Addition of positive integer to negative integer gives positive integer, zero
Addition of negative	3 + (-1) = 2	(-3) + 1 = (-2)	or negative integer.
integer to positive	3 + (-2) = 1 3 + (-3) = 0	(-3) + 0 = (-3) $(-3) + (-1) = (-4)$	
integer gives negative integer, zero	3 + (-4) = (-1) 3 + (-5) = (-2)	(-3) + (-2) = (-5) (-3) + (-3) = (-6)	negative integer to negative integer gives
or positive integer.	3 + (-6) = (-3)	(-3) + (-4) = (-7)	negative integer.

• Let us perform the addition and subtraction of integers without the help of number line. Let us understand the following :

Explanation 1: If a negative integer comes after the sign of addition or subtraction, remove the negative sign and change the sign of the process. e.g., 5 + (-3) = 5 - 3 = 2 and 5 - (-3) = 5 + 3 = 8

Explanation 2 : If a bigger number is to be subtracted from a smaller one, subtract the smaller number from the bigger one but the result should be given a negative sign.

e.g., 3 - 5 = (-2), 7 - 10 = (-3)


Explanation 3 : If the first number is negative and the process is of addition, according to the rule of commutative property, their order can be changed.

e.g.,
$$(-5) + 3 = 3 + (-5) = 3 - 5 = (-2)$$

 $(-7) + 9 = 9 + (-7) = 9 - 7 = 2$

Explanation 4: If the first number is negative and the process is of subtraction give negative sign to the sum of both the numbers. e.g., (-5) - 3 = (-8), (-7) - 9 = (-16)






Note : There is no need to write details shown in the brackets while solving the examples.




1. Write the absolute value :

(1) 17 (2)
$$(-18)$$
 (3) 0 (4) (-25) (5) 16

2. Make appropriate pairs :



Downloaded from https:// www.studiestoday.com



From the above two Tables, it can be said that the multiplication of zero with any positive or negative integer gives zero and zero multiplied by zero gives the result zero.

Thus, the multiplication of any integer with zero result into zero only.



Downloaded from https:// www.studiestoday.com

1411111	Jakatakakak hu	
	2 : Integers	
		and the second of the second

Look, understand and calculate :

(1)	$5 \times (-5) = (-25)$	(7)	$4 \times (-3) = \dots$
(2)	$(-12) \times 5 = (-60)$	(8)	$(-10) \times 4 = \dots$
(3)	$0 \times 12 = 0$	(9)	0 × 8 =
(4)	$(-5) \times (-10) = 50$	(10)	$(-4) \times (-5) = \dots$
(5)	$(-120) \times 1 = (-120)$	(11)	35 × 1 =
(6)	$1 \times 45 = 45$	(12)	$1 \times (-14) = \dots$

Properties of multiplication for integers :

In Standard 5, we have studied about the properties of multiplication of whole numbers. Now in the same way, we will study about the properties of multiplication of integers :

1. Look, understand and complete :

(-5)	×	11	=	(-55)
Integer		Integer		Integer
(-10)	×	(-15)	=	150
Integer		Integer		Integer
•••••	×	(-7)	=	0
Integer		Integer		Integer
5	×	(-7)	=	
Integer		Integer		Integer

It is clear from the above Table that the product of any two integers will be an integer only.

2. Look, understand and complete :

 $(-7) \times 10 = 10 \times (-7) = (-70)$ $(-3) \times (-7) = (-7) \times (-3) = 21$ $8 \times 0 = 0 \times 8 = \dots$ $7 \times (-6) = (-6) \times \dots = (-42)$



Downloaded from https:// www.studiestoday.com



It is clear from the above Table that the multiplication of any two integers can be done in any order. The result will be the same.

 3. Look and understand the following :

 $[(-3) \times 4] \times (-5)$ or
 $(-3) \times [4 \times (-5)]$ or
 $[(-3) \times (-5)] \times 4$
 $= (-12) \times (-5)$ $= (-3) \times (-20)$ $= (15) \times 4$

 = 60 = 60 = 60

 Complete as above :

 $[(-4) \times 5] \times (-7)$ or
 $(-4) \times [5 \times (-7)]$ or
 $[(-4) \times (-7)] \times 5$
 $= (-20) \times \dots = (-4) \times \dots = 140$ = 140

Thus, for the multiplication of any three integers taking a group of first two integers and then multiplying the product by the third integer, the result will be the same.

If we multiply any integer by 1, then we get the same integer as the result. Thus, 1 is called neutral (identity) number for multiplication.

Distribution of multiplication over addition :

We know that multiplication is repeated addition.


Look and understand :


$$(-4) + (-4) + (-4) + (-4) + (-4) = (-4) \times 5 = (-20)$$

$$(-4) + (-$$



Downloaded from https:// www.studiestoday.com






3. Number square - use the horizontal and vertical keys and fill the number-square :

	Horizontal key		Vertical key
А	The smallest whole no. of two digits	А	$(-21) \times (-5)$
В	34 + (-4)	D	Absolute value of (-124)
С	18 × 3	F	$(-9) \times 7$
Е	$(-23) \times (-21)$	G	21 + 14
F	(-8) + (-57)	Ι	The largest whole number of
G	13×30		four digits.
Н	(-6) - (-200)		27 - (-27)
Ι	$(-19) \times (-5)$	Κ	(-6) + 28
L	$7 \times (-6)$		



Downloaded from https:// www.studiestoday.com



Downloaded from https:// www.studiestoday.com

Square and Square Root


• Let us remember :

You have already learnt about square and its area. Measure of all the sides of square are equal, aren't they ? Area of a Square = Length \times Length. Let us do some activity with the help of a square having measure of 1 cm length to solve the following questions :

	Questions	Figure	Area (sq. cm)
•	What is the area of a square with length 1 cm?		$1 \times 1 = 1$
•	What is the area of a square with length 2 cm ?		$2 \times 2 = 4$
•	What is the area of a square with length 3 cm ?	Image: Constraint of the second se	$3 \times 3 = 9$
•	What is the area of a square with length 4 cm?	Image: select	
•	What is the area of a square with length 5 cm?	Image: select	

The area of a square having length 1 to 5 cm will be 1, 4, 9, 16, 25 sq cm respectively.





1, 4, 9 and 16 are squares of 1, 2, 3 and 4 respectively. These numbers are called perfect squares.

If the given number is a square of a number, it is called a perfect square number.

Therefore, $9 \times 9 = 9^2 = 81$

 $10 \times 10 = 10^2 = 100$

Here, 81 and 100 are perfect squares.

- Every number is not a perfect square number.
- e.g., 12 is not a square of any number, therefore 12 is not a perfect square number.

Think : How many such perfect squares can be found out ?



3 : Square and Square Root

Table : 1

Complete the following Table :

Number	Square	Perfect square number	Number	Square	Perfect square number	
1	$1 \times 1 = 1^2 = 1$	1	11			
2	$2 \times 2 = 2^2 = 4$	4	12			
3		9	13	$13 \times 13 = 13^2 = 169$	169	
4			14	•••••		
5			15			
6			16	$16 \times 16 = 16^2 = 256$	256	
7			17	•••••		
8	$8 \times 8 = 8^2 = 64$	64	18	$18 \times 18 = 18^2 = 324$	324	
9			19			
10	$10 \times 10 = 10^2 = 100$	100	20			

Based on the above Table 1, answer the following questions :

• Which digits are at the unit place in a perfect square ?

• Which digits do not come at unit place in a perfect square ?

- Answer :
- A perfect square has either 0, 1, 4, 5, 6 or 9 at its unit place.

• Digits other than 0, 1, 4, 5, 6 or 9 at the unit place of a number cannot be a perfect square. That means, the digits 2, 3, 7 or 8 never come at the unit place of a perfect square.





Example 2: Do the below given numbers have the possibility of being a perfect square ? Verify it.

(1) 5287 (2) 4302 (3) 361 (4) 3648 (5) 25 (6) 256

Solution :

- (1) The given number is not a perfect square because 7 is at its unit place.
- (2) The given number is not a perfect square because 2 is at its unit place.
- (3) The given number can be a perfect square because there is 1 at the unit place.
- (4) The given number is not a perfect square because its unit place is 8.
- (5) The given number can be a perfect square because its unit place is 5.
- (6) The given number can be a perfect square because its unit place is 6.
- **Example 3**: Write two such numbers looking at their unit place, it can be concluded that the numbers are not perfect squares.

Solution : (1) 3412 (2) 5007

MATHEMATICS



- 1. Is your roll number a perfect square number ? Why ?
- 2. In 1 to 100, how many perfect squares are found ? Note them down.
- 3. Which of the following numbers has the possibility of having perfect squares :
 - (1) 9285 (2) 2312 (3) 2307 (4) 2001 (5) 2305
 - (6) 2508
 (7) 2160
 (8) 44101
 (9) 1069
- **4.** Write ten such numbers, looking at their unit place, it can be concluded that they are not perfect squares.
- Write the squares of given numbers and complete the Table :

Table : 2

Number	Square	Number	Square	Number	Square
21	$21 \times 21 = 441$	26	$26 \times 26 = 676$	35	
22		27		40	$40 \times 40 = 1600$
23		28		50	
24	$24 \times 24 = 576$	29	$29 \times 29 = 841$	100	
25		30		500	500×500=250000

STD.7





Based on Table 1 and Table 2, complete Table 3 and Table 4 and get to the conclusion :

		Table : 3		
Unit place of perfect square numbers	1	4	6	9
	1	4	16	9
	81	64	36	49
Perfect square	121	144	196	169
numbers		•••••	•••••	
	••••••			
	••••	••••	•••••	•••••
Unit place digit of the	1 or 9		4 or 6	3 or 7
original number				
	The number having 1 or 9 at its unit	The number having 2 or 8 at its unit		The number having 3 or 7 at its unit
Conclusion	place will have 1 at	place will have 4 at	•••••	place will have 9 at
	the unit place of	the unit place of		the unit place of
	perfect square of that number.	perfect square of that number.		perfect square of that number.

Table : 4

The last digit of the perfect may square number	25	00 I	0000
	25	100	10000
Perfect square	225	400	
numbers 👘			
	••••		
Unit place digit of the	5	0	00
original number			(last two digits)
Conclusion	The number with 5 at its unit place will surely have 25 as the last two digits.	The numbers with 0 at its unit place will surely have 00 as the last two digits.	The number with 0 at its unit and ten's place, will surely have 0000 as the last four digits.





Example 4 : By squaring the below given number, which digits are found at the unit place.

Number	Digit found at the unit place by squaring the number	Number	Digit found at the unit place by squaring the number
225	5	53004	6
157	9	83091	1

Example 5 : Mention the last two digits of the square of the following number : (1) 25 (2) 30 (3) 35

Solution : (1) We get 25. (2) We get two zeros. (3) We get 25.

Example 6 : How many zeros are obtained by squaring the below given numbers ? (1) 20 (2) 200

Solution : (1) We will get two zeros because zero is only at the unit place.

(2) We will get four zeros because it has zero at its unit and ten's place.

Based on the above conclusion answer the following

- (1) By squaring the total number of students in your class, what will be the digit at unit place ?
- (2) If you square your roll number which digit will be obtained at the unit place ?
- (3) If you square the number of your birthdate which digit will be obtained at the unit place ?
- (4) If you square the number of your weight, which digit will be obtained at its unit place ?
- (5) If you square the number of pages of your notebook, which digit will be obtained at its unit place ?

Method to obtain square of a number having 5 at its unit place :

• Study the following Table :

MATHEMATICS


Number	Square	In perfect squares, digit formed at hundredth and thousandth place
15	225	$2 = 1 \times 2$
25	625	$6 = 2 \times 3$
35	1225	$12 = 3 \times 4$
45	2025	$20 = 4 \times 5$
55	3025	$30 = 5 \times 6$

STD.7





By studying the above Table we conclude that if we square a number with 5 in its unit place, the perfect square formed will have 2 and 5 i.e. (25) at tens and unit place respectively. The number preceding 25 is the product of two consecutive numbers. For example, to get square of 35, first multiply the number 3 in tens place with the number 4 next to it. Write 25 at the end of the result. So, $35^2 = 1225$



By squaring a number having 5 in its unit's place, the number formed by tens and unit place is always 25.



Write the squares of the following numbers :

(1) $75^2 = \dots$ (2) $65^2 = \dots$ (3) $85^2 = \dots$ (4) $105^2 = \dots$

Square root :

(1) $1 \times 1 = 1^2 = 1$	(2)	$4 \times 4 = 4^2 = 16$
(3) $10 \times 10 = 10^2 = 100$	(4)	$15 \times 15 = 15^2 = 225$

Here, 1, 16, 100, 225 are the perfect squares obtained by squaring 1, 4, 10 and 15 respectively. 16 is the square of 4, therefore 4 is the square root of 16. 100 is the square of 10, therefore 10 is the square root of 100.





Complete the following Table :

Square	Square root	Square	Square root
$11^2 = 121$	$\sqrt{121} = 11$	$16^2 = \dots$	
$12^2 = 144$	$\sqrt{144} = 12$	$17^2 = 289$	$\sqrt{289} = 17$
$13^2 = \dots$	$\sqrt{169} = 13$	$18^2 = 324$	
$14^2 = \dots$		$19^2 = \dots$	$\sqrt{361} = 19$
$15^2 = \dots$		$20^2 = \dots$	

Finding the square root by the method of indivisible factors :

Example 6 : Find the square root of 36.

Solution :	2	36	$36 = \underline{2 \times 2} \times \underline{3 \times 3}$
	2	18	$= 2^2 \times 3^2$
	3	9	$= (2 \times 3)^2$
	3	3	$= 6^2$
		1	$\therefore \sqrt{36} = \sqrt{6^2} = 6$
			\cdots $\sqrt{30} = \sqrt{0} = 0$

Example 7 : Find the square root of 1089.

Solution :	3	1089	$1089 = \underline{3 \times 3} \times \underline{11 \times 11}$
	3	363	$= 3^2 \times 11^2$
	11	121	$= (3 \times 11)^2$
	11	11	$= 33^2$
		1	$\therefore \sqrt{1089} = 33$


• Do it yourself : Find the square root of 25, 64 and 100.

Indivisible factors of $32 = 2 \times 2 \times 2 \times 2 \times 2$

Here, 2 has total two pairs and one 2 remains alone, therefore 32 is not a perfect square.



Downloaded from https:// www.studiestoday.com



(1) 64	(2) 100	(3) 484	(4) 900	(5) 1150
(6) 3136	(7) 1225	(8) 1764	(8) 12100	(10) 3249

2. Are 42 and 50 the perfect squares ? Why ?

Example 8 : Is 252 a perfect square ? By which smallest number should it be multiplied, so that the result becomes a perfect square ?

Solution :	2	252	$252 = 2 \times 2 \times 3 \times 3 \times 7$
	2	126	In factors of 252 pair of 7 is not formed, so 252 is not a
	3	63	perfect square. If we multiply both the sides by 3, we get
	3	21	perfect square.
	7	7	$252 \times 7 = \underline{2 \times 2} \times \underline{3 \times 3} \times \underline{7 \times 7}$
		1	By multiplying 252 by 7, we get perfect square.


Example 9: Is 1620 a perfect square ? By which smallest number should it be divided so that the result becomes a perfect square ?

2	1620								
2	810	$1620 = \underline{2 \times 2} \times \underline{3 \times 3} \times \underline{3 \times 3} \times 5$							
3	405	In factors of 1620 pair of 5 is not formed, so 1620 is							
3	135	not a perfect square.							
3	45	If we remove factor 5 from both the sides i.e. by dividing							
3	15	by 5 on both the sides we get perfect square.							
5	5	$1620 \div 5 = 2 \times 2 \times 3 \times 3 \times 3 \times 3 \times 5 \div 5$							
	1								
$= 2 \times 2 \times 3 \times 3 \times 3 \times 3$									

Now every indivisible factor forms a pair.

So, by dividing 1620 by 5, we get a perfect square.





Square and square root of a fraction :

We have learnt to find the squares of an integer. To find the square of a fraction, we place square of numerator in numerator's place and the square of denominator in denominator's place.


$$\left(\frac{1}{2}\right)^2 = \frac{1}{2} \times \frac{1}{2} = \frac{1 \times 1}{2 \times 2} = \frac{1}{4}$$
$$\left(\frac{2}{5}\right)^2 = \frac{2}{5} \times \frac{2}{5} = \frac{2 \times 2}{5 \times 5} = \frac{4}{25}$$
$$\left(\frac{3}{4}\right)^2 = \frac{3}{4} \times \frac{3}{4} = \frac{3 \times 3}{4 \times 4} = \frac{9}{16}$$

In the same way, square root of a fraction means placing the square root of numerator at numerator's place and square root of denominator at denominator's place.

$$\left(\frac{1}{2}\right)^2 = \frac{1}{4}, \text{ thus } \sqrt{\frac{1}{4}} = \sqrt{\frac{1^2}{2^2}} = \frac{1}{2}$$
$$\left(\frac{3}{4}\right)^2 = \frac{9}{16}, \text{ thus } \sqrt{\frac{9}{16}} = \sqrt{\frac{3^2}{4^2}} = \frac{3}{4}$$
$$(1.5)^2 = \left(\frac{15}{10}\right)^2 = \frac{225}{100}, \text{ so, } \sqrt{2.25} = \sqrt{\frac{225}{100}} = \sqrt{\frac{15^2}{10^2}} = \frac{15}{10} = 1.5$$

Example 10 : Find the value of $\sqrt{\frac{25}{49}}$.

Solution : Now, we factorize the numerator and denominator.




Example 11 : Find the square root of $11\frac{14}{25}$.

Solution : Here, $11\frac{14}{25}$ is a vulgar fraction. Let us convert it into improper fraction and then find the square root by the method in the above example.

$$11\frac{14}{25} = \frac{289}{25}$$




Downloaded from https:// www.studiestoday.com



Example 12 : Find the square of 12.25.


Solution : Here 12.25 is a decimal fraction, so it can be transformed into simplefraction and square root could be found out.512252100




1. Write the digit in unit place by squaring the number :

Sr. No.	Number	Digit in unit place by squaring the number
1	125	
2	137	
3	140	
4	78	
5	95	
6	108	





Downloaded from https:// www.studiestoday.com





Downloaded from https:// www.studiestoday.com



Let us remember :

4

- The amount at which a trader buys a product is known as its Cost price.
- After buying a product, the additional expenditure incurred on a product is called Additional expense. The expenditure on labour, rent, octroi tax, maintenance work is called additional expense.
- The sum of cost price and additional expense is called Net price. When there is no extra expenditure, the cost price is considered the Net price.

```
Net price = Cost price + Additional expense
```

- The amount at which a trader sells the product is called the Sale price.
- The excess amount got over the cost price after selling the product is called the **profit**.

Profit = Sale price - Net price

So, Profit = S.P. - N.P.

The amount got at the time of sale less than the net amount is called the Loss.
 Loss = Net price - Sale price

So, Loss = N.P. - S.P.

When there is profit, Sale price = Net price + Profit

When there is loss, Sale price = Net price - Loss

Now, find the answers from above information :


A trader bought a TV for ₹ 9950. He paid ₹ 50 as labour charge to bring the TV home. Selling the TV at ₹ 10,700, he earned the profit of ₹ 700.

Cost price	=₹	Additional expense = ₹		
Net price	=₹	Sale Price	=₹	
Profit	=₹	Profit	= %	

Let's learn something new :

• In Std. 6, we learnt how to find profit or loss in terms of percentage. Now on the basis of profit and loss percentage and net price, let's understand how to find the selling price.

MATHEMATICS 33 STD. 7



1. Fill the following Table after calculation :

Sr. No.	Cost price (in ₹)	Addi. Exp. (in ₹)	Profit (in %)	Loss (in %)	Sale price
1.	60	(m () _	5	(III / 0) _	
2.	40		10	_	
3.	1000	_	12	_	
4.	240	_	_	15	
5.	1500	_	_	5	
6.	24	_	-	12.5	
7.	1650	150	—	5	•••••
8.	750	50	—	10.5	·····
9.	3800	200	15.5	_	••••

- Vairagi bought a table for ₹ 450. There is ₹ 50 additional expenses on this table.
 For earning 12 % profit, what should be the sale price ?
- Vivekbhai bought a TV for ₹ 16,000. ₹ 200 was spent on transportation and labour. For earning 12 % profit, what price should it be sold ?



Downloaded from https:// www.studiestoday.com

4 : Profit-Loss

- 4. Deepali has bought a sofa set for ₹ 14,500. There is an additional expense of ₹ 500 on it. Since she did not like the sofa set, she sold it bearing a loss of 8.5 %. For how much did she sell the sofa set ?
- 5. Akhileshbhai bought a house for ₹ 9,50,000. ₹ 50,000 was spent on colour. For earning a profit of 14.5 %, how much should sell the home for ?
- 6. Tamiza sold off her Activa costing ₹ 55,000 to Sharmin earning a profit of 14 %. How much amount should Tamiza have got ?
- Brokerage, Commission :

Nareshbhai wants to buy an old tractor. He tells this to Hareshbhai.

Hareshbhai : I remember that there is a tractor used for two years only.

Nareshbhai : Wow, that is very nice.

Hareshbhai : I will show you a tractor as you wish to buy it at an economical price. But, if you buy that tractor, you should give me ₹ 2000.

Nareshbhai : Ok. Kindly show me that tractor.

Hareshbhai : Come on (Going to Pareshbhai's field).

Pareshbhai... Come on this side. Look, my friend Nareshbhai had come. He wants to buy your tractor, if he likes it. Talk to him on your own, but if it is sold, your must give me ₹ 1000.

On seeing the tractor, Nareshbhai likes it. He asks the cost and at the end of negotiation, Nareshbhai buys the tractor on giving $\overline{\mathbf{x}}$ 1,80,000. According to the condition gives $\overline{\mathbf{x}}$ 2000 to Hareshbhai. Pareshbhai calls Hareshbhai and gives $\overline{\mathbf{x}}$ 1000 as decided earlier for helping him to see the tractor.

Now answer the following :

- (1) At what price did Pareshbhai sell the tractor ?
- (2) At what price did Nareshbhai buy the tractor ?
- (3) How much money did Pareshbhai give to Hareshbhai? Why?
- (4) How much money did Nareshbhai give to Hareshbhai ? Why ?
- (5) How much money did Pareshbhai actually get on selling the tractor ?
- (6) How much money was actually paid by Nareshbhai to buy the tractor ?
- (7) From whom did Hareshbhai get money ? How much ?
- (8) Hareshbhai did not actually buy or sell any thing, yet he get money, why ?

A person who bring the buyer and seller together is called a broker.

In the above conversation, Hareshbhai is a broker.



4 : Profit-Loss

When any product is to be sold or bought, the amount that a broker gets from either seller or buyer or both is called brokerage. In the above conversation Hareshbhai takes ₹ 2000 from Nareshbhai and ₹ 1000 from Pareshbhai as brokerage.

Brokerage is a kind of additional expense.

Amount received by the seller = S.P. - Brokerage Amount to be paid by the buyer = C.P. + Brokerage

Example 2 : Prakruti bought an old sofa set for ₹ 9600, with the help of a broker at 2 % brokerage. At what price did she get the sofa set ?

Solution : 2 % brokerage means on the cost price ₹ 100 the brokerage is ₹ 2.

:. Here, brokerage = 2% of ₹ 9600

$$= \not\in \left(9600 \times \frac{2}{100}\right) = \not\in 192$$

 \therefore Net price = C.P. + Brokerage

- ∴ Prakruti got the sofaset for ₹ 9792.
- Example 3 : Kasambhai bought a shop for ₹ 6,00,000, through a broker. The broker took 1 % brokerage from the seller and 2 % brokerage from the buyer. How much brokerage did the broker get ?

Solution : Method 1 : The broker takes 1 % brokerage from the seller :

... Brokerage paid by the seller

= ₹ 6,00,000 + 1 %
= ₹
$$(600000 \times \frac{1}{100})$$

= ₹ 6000

The broker takes 2 % brokerage from the buyer :

... Brokerage paid by the buyer

= 2 % of ₹ 6,00,000
= ₹
$$(600000 \times \frac{2}{100})$$

= ₹ 12,000



4 : Profit-Loss

:. Total brokerage earned by the broker = Brokerage paid by the seller +

Brokerage paid by the buyer

= ₹ 6,000 + ₹ 12,000 = ₹ 18,000

∴ The broker gets total ₹ 18,000 as brokerage.

Method 2 : The brokerage earned by the broker = 2% + 1%

= 3 %

In this way, total brokerage can be calculated at the rate of 3 % on ₹ 6,00,000.

Total brokerage earned by the broker = \neq (600000× $\frac{3}{100}$)

= ₹ 18,000



- 1. By paying 2 % brokerage for buying an old scooter for ₹ 15,000 through a broker. At what amount is the scooter bought ?
- 2. A car was sold at ₹ 80,000 through a broker. The broker charged 2.5 % brokerage from both the seller and the buyer. What is the total amount of brokerage received by the broker ?
- 3. Vinodbhai sold his shop through a broker for ₹ 7,50,000. The broker charged 1 % brokerage for this work. What amount did Vinodbhai get on selling the shop ?
- 4. Dharmendrabhai sold a plot through a broker for ₹ 8,80,000. The broker charged 1.5 % brokerage for this work. What amount did Dharmendrabhai get on selling the plot ?
- 5. Manojbhai bought an old tractor through a broker for ₹ 2,50,000. The broker charged 1 % brokerage from the seller and 2 % brokerage from the buyer. What is the total amount of brokerage received by the broker ?
- Rebate (Discount) :


MRP is the maximum amount at which a product can be sold. **MRP** (Maximum Retail Price) is printed on the product.

When a trader takes an amount less than the printed amount, that much amount is called **Rebate** for the buyer.

Rebate = Printed price \times % or rebate

Amount to be paid = Printed price - Rebate





	4 : Profit-Loss					
N	0.	Product	MRP (₹)	% rebate	Amount of the rebate (₹)	Amount to be paid (₹)
(1	l)	Towel				
(2	2)	Curtain				
(3	3)	1 kg Biscuits				

2. Calculate and fill the Table :

No.	Book	MRP (₹)	Rebate	Amount to be paid (₹)
(1)	My Experiments With Truth	20	10 %	
(2)	Srimad Bhagvad Geeta	65	20 %	
(3)	Bible	60	20 %	
(4)	Day Dreams	80	5 %	
(5)	Panchtantra	120	15 %	

- 3. **Project** : From the newspaper, cut the advertisement showing rebate and find the cost of that product.
- Practical sums :
- Example 6 : A saree trader buys some sarees at a rebate of 20 % on the printed price of ₹ 600. If he sells these sarees at ₹ 540 each, how much profit or loss will he make per saree ? How much rebate in percentage does a customer get on each saree ? Rebate for the trader = Printed price × % of rebate

= ₹ (600 × 20 %)= ₹ $(600 × \frac{20}{100})$ = ₹ 120 Net price of the saree = Printed price - Rebate = ₹ 600 - ₹ 120 = ₹ 480



Downloaded from https:// www.studiestoday.com

4 : Profit-Loss

The trader sells this saree for ₹ 540.

(1) Profit earned per saree = Selling price - Net price

Rebate earned by the customer = Printed price - Price paid

- ∴ Rebate earned by the customer on ₹ 600 = ₹ 60
- (2) % rebate earned by the customer = $\mathbf{E}\left(100 \times \frac{60}{600}\right) = 10 \%$
- \therefore The customer earned a rebate of 10 % on each saree.




- A trader buys some T-shirts at 10 % rebate on printed price ₹ 450 per piece.
 For a long time he could not sell any T-shirt, so he sells it for ₹ 360 per piece.
 What amount of profit or loss per piece does he make ? How much percentage of rebate does a customer earn ?
- 2. A trader buys some TVs at 20 % rebate on printed price ₹ 18,000. For a long time he could not sell any TV. So he sells it by giving a rebate of ₹ 2700 on printed price, what amount of profit or loss does he make ? How much percentage of rebate does a customer earn ?




1. Calculate the sale price on the basis of the given information :

Cost price (₹)	Additional expense (₹)	Profit (%)	Loss (%)	Sale price (₹)
1800	_	_	7	
630	70	10	_	
1050	150	4	_	





- A trader buys some cameras with printed price ₹ 750. If he sells each camera giving 10 % rebate, how much money could a customer have to pay ?
- 3. A saree trader brings some sarees at 25 % rebate on ₹ 1600 printed price. He sells each saree at ₹ 1440 per piece. What amount of profit or loss does he make ? How much percentage of rebate does a customer earn ?
- **4.** A television costs ₹ 14,750. If a customer gets a rebate of 20 % on that, at what amount does the customer get the television ?
- 5. 10 % rebate is given in a T-shirt shop. Sohan buys 5 T-shirts with ₹ 300 printed price on each. He sells all the T-shirts for ₹ 1485. Does he earn profit or suffer loss ? How much amount ?



Practice 1

1.	(1) ₹ 63	(2) ₹ 44	(3) ₹ 1120	(4) ₹ 204	
	(5) ₹ 1425	(6) ₹ 21	(7) ₹ 1710	(8) ₹ 716 (9)	₹ 4620
2.	₹ 560 3. ₹	18,144 4 .	₹ 13,725 5. ₹	[*] 8,55,000 6. ₹ 6	52,700

Practice 2

1. ₹ 15,300 **2.** ₹ 4000 **3.** ₹ 7,42,500 **4.** ₹ 8,66,800 **5.** ₹ 7500

Practice 3

1. (1) ₹ 18.75, ₹ 106.25(2) ₹ 75, ₹ 675(3) ₹ 2, ₹ 382. (1) ₹ 18(2) ₹ 52(3) ₹ 48(4) ₹ 76(5) ₹ 102

Practice 4

1. (1) Loss ₹ 45, 20 % Rebate (2) Profit ₹ 900, 15 % Rebate

Exercise

- 1. (1) ₹ 1674 (2) ₹ 770 (3) ₹ 1248
- **2.** (1) ₹ 675 **3.** (1) ₹ 240 Profit, 10 % Rebate
- **4.** ₹ 11,800 **5.** ₹ 135 Profit

MATHEMATICS

Downloaded from https:// www.studiestoday.com

STD.7

4 : Profit-Loss

Let us understand

The value of a product increases at every step from production to distribution. VAT means Value Added Tax. VAT is such a tax that is charged on the increase of value of the product at every step. In fact VAT is a kind of Sales Tax charged on the sale price of a product.

According to the rules of VAT, a trader has to pay VAT to the seller at the time of buying any product. In the same way VAT is charged from the buyer at the time of selling the product. After this, at the time of calculating the VAT amount to be paid to the government, the trader pays the difference of the amount paid by the trader to the government and the amount charged by the trader.

Example : A trader sells some product at ₹ 60,000 bought at the price of ₹ 50,000.

If VAT is 4 %

(1) Find the amount of VAT paid to the government by the trader ?


(2) Find the amount of VAT charged by the trader from the customer ?

(3) Lastly, find the amount of VAT to be paid by the trader to the government. Solution :

(1) The amount of VAT paid to the government by the trader = 4 % of ₹ 50,000

= ₹ 2000

- (2) The amount of VAT charged by the trader from the customer = 4 % of ₹ 60,000 = ₹ 2400
- (3) Lastly, the amount of VAT to be paid to the government by the trader
 = VAT charged at the time of selling VAT charged at the time of buying
 = ₹ 2400 ₹ 2000 = ₹ 400



Parallel Lines

• Let us remember :

5

Sr.	Measure of the given angle	Measure of the complementary angle	Measure of the supplementary angle	Measure of the angles of a linear pair	Measure of the vertically opposite angles
1.	75°	15°	105°	105°	75 ⁰
2.	80°		100 ^o		
3.	60°			120°	
4.	72°	18 ⁰			
5.	15°				15°

Fill in the blanks after studying the following figure

- (1) $m \angle AOC = \dots$
- (2) $m \angle \text{COB} = \dots$
- (3) $m \angle BOD = \dots$
- (4) Measure of the complementary angle of $m \angle AOD$ is =
- (5) Measure of the supplementary angle of complementary angle of $m\angle COB =$

70⁰


D


0

B


C


Let us learn new :





Two opposite sides of a blackboard






What kind of distance is there between the opposite sides in all the above figures ?


Activity 1 :


- Take 1 meter long string.
- Join the two ends and make a loop.
- With the help of two pencils, keep the string stretched up.
- Measure the distance between the two straight lines with the help of scale.

Questions :

- (1) Do \overline{AB} and \overline{CD} intersect each other ?
- (2) Think and say that does a straight line form the sides of a scale ?
- (3) Is the distance between both the lines same ?
- (4) Measure the distance between the two lines, with the help of a scale.
- (5) By extending both lines, will it intersect at any one place ?
- Study the figures given here and answer the questions based on them :







- 1. In figure (1), (2) and (3), draw the figure by extending lines l and m and name the point of intersection.
- 2. By extending lines l and m in figure (4), will they intersect each other ? Why ?

In a plane, two distinct lines which do not intersect and the distance between the lines is same, such lines are known as parallel lines.

In figure (4) line l and line m are parellel. Symbolically, it is written as $l \parallel m$ and read as line l is parallel to line m.

- Activity 2 : Find out the examples of such things found in our day to day life having opposite sides parallel and write their names :

The distance between parallel lines should be found out with the help of set squares :

Activity 3 :


- Take one page from a notebook.
- Measure the distance between the opposite sides with the help of a scale.
- Note down the distance everytime.
- Is the distance same everytime ?
- Think the opposite sides of the page are parallel or not ?
- Activity 4 :
 - Draw two lines which includes the two sides of a scale.
 - Name them as *l* and *m*.
 - Take a point A on line *m*.

MATHEMATICS

Downloaded from https:// www.studiestoday.com

45

STD.7



Study figures (5), (6) and (7), given above and answer the questions given below :

(1) In figure (5), at which point does line l intersect the other two lines ?

- (2) In figure (6), at which points, does line l intersect the other two lines ?
- (3) In which figure, the point of intersection is same for all the three lines ?
- (4) For figure (7), give the answers to the details of the following table.

Line	Which two lines are intersected ?	Point of intersection
l	m and n	C and A
m		
n		

STD.7



MATHEMATICS

5 : Parallel Lines

In figure (6) and (7) line l intersects line n and line m in two different points. Therefore, line l is called the transversal of line m and line n. While in figure (5) line l is not the transversal of line m and line n. In figure (7) each line is the transversal of the other two lines.

Transversal : If a line intersects two other lines in the same plane at two distinct points, it is called a transversal of the two lines.


• Angles formed by a transversal :

• Activity 5 :

- Take three straws.
- Arrange three straws with the help of pins as shown in figure (6).
- Now, think about the angles formed by them.
- Think about the point of intersection.
- Give them names.


Study the figure (8) given here and answer the questions given below :

- (1) In all, how many lines are there ?
- (2) Which line is transversal of other lines and at which points does it intersect other lines ?
- (3) In all, how many angles are formed ? Name them,
- (4) How many angles are formed in the interior of \overrightarrow{AB} and \overrightarrow{CD} ? Name them,
- (5) How many angles are formed at the exterior, of \overrightarrow{AB} and \overrightarrow{CD} ? Name them,
- Types of angles formed by a transversal :
 Look and understand them :



 $\angle 1$, $\angle 2$, $\angle 3$ and $\angle 4$ are interior angles (9)

MATHEMATICS



0

(8)

STD.7

D

 $\angle 5$, $\angle 6$, $\angle 7$ and $\angle 8$ are exterior angles (10)


5 : Parallel Lines

	Number of angles	Angles
Interior angles of the two lines (fig. (9))1		
Exterior angles of the two lines (fig. (10))		

• A pair of alternate angles :

In figure (11), $\angle 3$ and $\angle 6$ are a pair of alternate angles. In the same way $\angle 4$ and $\angle 5$ are a pair of alternate angles.

Thus, we have two pairs of alternate angles.



3

D

(12)

STD.7

Figure No.	Angles	Pair of Alternate angles
11	$\angle 3$ and $\angle 6$	$\angle APQ$ and $\angle PQD$
	$\angle 4$ and $\angle 5$	\angle and \angle


There are two pairs of alternate angles.

Corresponding angles :

MATHEMATICS

In figure (12), $\angle 1$ and $\angle 5$

- $\angle 2$ and $\angle 6$
- $\angle 3$ and $\angle 7$



Thus, we have four pairs of corresponding angles.

Figure No.	Angles	Pair of Corresponding angles
(12)	$\angle 1$ and $\angle 5$	$\angle XPA$ and $\angle PQC$
	$\angle 2$ and $\angle 6$	\angle and \angle
	$\angle 3$ and $\angle 7$	∠ and ∠
	$\angle 4$ and $\angle 8$	∠ and ∠

There are four pairs of corresponding angles.




Interior angles on the same side of a transversal :

In the adjoining figure (13)

 $\angle 4$ and $\angle 6$

 $\angle 3$ and $\angle 5$ are the interior angles on the

same side of the transversal.



Thus, there are two pairs of interior angles on the same side of the transversal.


Fig. No.	Angles	Name of angles
(13)	$\angle 4$ and $\angle 6$	$\angle BPQ$ and $\angle PQY$
	$\angle 3$ and $\angle 5$	\angle and \angle

There are two pairs of interior angles on the same side of the transversal.


• Taking the help of your teacher prepare a Power-Point presentation and observe alternate angles, corresponding angles and angles on the same side of the transversal.



1. In the figures given below, write which lines form a pair of parallel lines and write them in the form of symbols :



Downloaded from https:// www.studiestoday.com



5 : Parallel Lines

Conclusion : (1) The measure of angles of the pair of alternate angles are

- (2) The measure of angles of the pair of corresponding angles are

For the angles formed by the transversal of two parallel lines :

- Measures of both angles of every pair of alternate angles are equal.
- Measures of both angles of every pair of corresponding angles are equal.
- The sum of the measure of both angles in every pair of interior angles on the same side of the transversal is 180°. (Both angles of each pair of interior angles on the same side of the transversal are supplementary).

Example 1 : In the figure given below $\overrightarrow{AB} \parallel \overrightarrow{XY}$ and \overrightarrow{CD} intersect \overrightarrow{AB} and \overrightarrow{XY} at point P and Q respectively.

A

 75°

STD.7

 $105^{\circ}Q^{75^{\circ}}$

D

B

1050

If $m \angle APC = 105^{\circ}$, find the measure of the

remaining angles.

Solution :

 $m \angle APC = 105^{\circ}$ $\therefore m \angle XQP = 105^{\circ}$ (corresponding angles) $m \angle XQP = 105^{\circ}$ $\therefore m \angle BPQ = 105^{\circ}$ (alternate angles) $m \angle BPQ = 105^{\circ}$ $\therefore m \angle YQD = 105^{\circ}$ (corresponding angles) Now, $m \angle BPQ + m \angle YQP = 180^{\circ}$ (linear pair as they are interior angles on the same side of the transversal)

 $105^{\circ} + m \angle YQP = 180^{\circ}$

$$\therefore m \angle YQP = 180^{\circ} - 105^{\circ}$$


 $m \angle YQP = 75^{\circ}$


MATHEMATICS

 $m \angle YQP = 75^{\circ}$ \therefore $m \angle APQ = 75^{\circ}$ (alternate angles)

 $m \angle APQ = 75^{\circ}$ \therefore $m \angle XQD = 75^{\circ}$ (corresponding angles)

 $m \angle YQP = 75^{\circ}$ \therefore $m \angle BPC = 75^{\circ}$ (corresponding angles)









1. Fill in the blanks :

- (1) Line t_1 and t_2 are parallel lines. We write in symbol as
- (2) In all angles are formed by a transversal of two lines.
- (3) pairs of alternate angles are formed by the transversal of two lines.
- (4) pairs of corresponding angles are formed by the transversal of two lines.
- 2. In the figure given here, state which of the line is the transversal of other two lines :



STD.7

- 3. $\overrightarrow{JK} \parallel \overrightarrow{MN}$. \overrightarrow{AB} is the transversal. \overrightarrow{AB} intersects \overrightarrow{JK} in point O and \overrightarrow{MN} in point B. Draw figure and write all the pairs of alternate angles, corresponding angles and interior angles on the same side of the transversal.
- 4. $\overrightarrow{XY} \parallel \overrightarrow{MN}$. \overrightarrow{PQ} is the transversal. \overrightarrow{PQ} intersects \overrightarrow{XY} at point A and \overrightarrow{MN} at point B. If $\angle NBP = 55^{\circ}$, find the measure of the remaining angles.
- 5. Draw a figure in which $\overrightarrow{AB} \parallel \overrightarrow{CD}$. \overrightarrow{XY} is a transversal. \overrightarrow{XY} intersects \overrightarrow{AB} at point P and \overrightarrow{CD} at point Q. If $\angle XPA = 120^{\circ}$, find the measure of $\angle PQD$ and $\angle BPQ$.
- 6. For the parallel lines and their transversal if the measure of one of the angles of the corresponding pair is 110°, then find the measures of all the alternate pairs of angles.




Practice 1

- 1. (1) $\overrightarrow{AB} \parallel \overrightarrow{CD}$ (2) No (3) No (4) $\overrightarrow{EP} \parallel \overrightarrow{FQ}$
- **2.** (1) Line l_3 is a transversal of l_1 and l_2 .


MATHEMATICS


Downloaded from https:// www.studiestoday.com

53



Downloaded from https:// www.studiestoday.com





Questions :


MATHEMATICS

- (1) What is the perimeter of the rectangle given in figure 1?
- (2) What did you do to find the perimeter of the rectangle ?
- (3) How can we find the perimeter of a square ?
- (4) What is the perimeter of the square given in figure 2?



55

STD.7










Adding 2 tops to 4 tops, there will be 6 tops. That is 4 tops + 2 tops = 6 tops If we use the symbol y for a top, 4y + 2y = 6y



۲





3 marbles + 4 tops can be written only as 3 marbles + 4 tops. But it cannot be written as 7 marbles or 7 tops. If we use the symbol x for a marble and y for a top, it can be written as 3x + 4y.





Here, 3x and 7x and 4y and 2y are like terms. Thus addition can be done for only like terms. 3x and 4y are unlike terms. Addition of unlike terms is denoted by putting the sign of addition in between them.

Example 1 : Add 2x, 3x and 4x.

Horizontal method	Vertical method
2x + 3x + 4x	2 <i>x</i>
= (2 + 3 + 4)x	+ 3x
= 9x	+ 4x
	9x

Example 2 : Add $5a^2b^2$ and $6a^2b^2$.

Horizontal method	Vertical method
$5a^2b^2 + 6a^2b^2$	$5a^{2}b^{2}$
$= (5 + 6) a^2 \times b^2$	$+ 6a^2b^2$
$= 11a^2b^2$	$11a^{2}b^{2}$

Example 3 : Add $2x^2$ and 3x.

Here, $2x^2$ and 3x are unlike terms. Therefore, the addition of the unlike terms can be written as

Addition = $2x^2 + 3x$ **Example 4** : Add 5*ab*, 3*ab* and $4a^2b^2$. = $5ab + 3ab + 4a^2b^2$ = $(5 + 3)ab + 4a^2b^2$ = $8ab + 4a^2b^2$




1. Make pairs of like terms from that given below : $2x^2$, $3x^2$, 5xy, $6x^2y^2$, $9x^2y^2$, 7xy

MATHEMATICS 57 STD. 7


Downloaded from https:// www.studiestoday.com



- 2. Find the like terms from the given terms and add them : 4xyz, $2x^2y^2$, $3x^2y^2$, $4x^2$, $9x^2y^2$, $18x^2$, 6xyz, $10x^2$, 7xyz
- 3. Calculate the perimeter of the following figures :



Subtraction of a monomial from a monomial :
 See and understand it :



Subtracting 2 marbles from 6 marbles 4 marbles remain.

That is 6 marbles - 2 marbles = 4 marbles

If we use the symbol x for a marble, then 6x - 2x = 4x



Here, subtracting 3 tops from 4 marbles can be written as follows :

4 marbles - 3 tops

If we use the symbol x for a marble and y for top, it can be written as 4x - 3y. Thus, subtraction can be done for only like terms. For unlike terms, subtraction is denoted by putting the sign of subtraction in between them.





We know that, the number opposite to the number which is to be subtracted is added in subtraction.

Subtract 3 from 5 :

3 is to be subtracted from 5 i.e. the opposite number of 3 that is (-3) is to be added. That is 5 + (-3) = 2

In the same way subtracting (-8) from (-6), the opposite number of (-8) that is 8 is to be added.

That is (-6) + 8 = 2

In the same way, to subtract one polynomial from other polynomial, the opposite polynomial of the polynomial which is to be subtracted is added to the given polynomial. To subtract 3x from 5x, the opposite polynomial of 3x i.e. (-3x) is to be added.

That is 5x + (-3x) = 5x - 3x.

Example 5 : Subtract 8xy from 15xy.

Horizontal method	Vertical method
15xy - 8xy	15 <i>xy</i>
$= 15 \times xy - 8 \times xy$	-8xy
= (15 - 8) xy	7 <i>xy</i>
=7xy	

Example 6 : Subtract $-9a^2b^2c^2$ from $16a^2b^2c^2$.

Vertical method
$\frac{16a^2b^2c^2}{+9a^2b^2c^2}}{25a^2b^2c^2}$


From example 6, it is understood that subtract is to add the opposite of a given number.

Example 7 : Subtract 7xy from $5x^2$.

Here $5x^2$ and 7xy are unlike terms, their subtraction can be written as :

 $= 5x^2 - 7xy$





(1) $4x^2$ and $-6xy^2$ (2) $6x^3$ and $-2x^3$ (3) 9xy and 5xy (4) $-7x^3$ and $-8x^3y$ The teacher should check the calculation of Q1 and Q2 alone by the students :

• Addition of a monomial with a binomial :

Example 8 : Add 6x and 3x + 7.

Horizontal method	Vertical method
(6x) + (3x + 7)	6 <i>x</i>
= <u>6x + 3x</u> + 7 (Taking like terms together)	+ 3x + 7
= (6 + 3)x + 7	9x + 7
= 9x + 7	

Example 9 : Add $(3a^2b^2 - 4)$ and $5a^2b^2$.


Horizontal method	Vertical method
$= (3a^2b^2 - 4) + (5a^2b^2)$	
$= 3a^2b^2 - 4 + 5a^2b^2$	
$= 3a^2b^2 + 5a^2b^2 - 4$	
$= (3 + 5) a^2 b^2 - 4$	
$= 8a^2b^2 - 4$	

Example 10 : Add $m^2 - 7$ and 25.

Horizontal method	Vertical method
	$m^2 - 7$
	+ 0 + 25
	$m^2 + 18$



Downloaded from https:// www.studiestoday.com



(4) -5n and 8n + 7 (5) $8x^2 + 7$ and $-8x^2$ (6) 3xy - 5 and 9xy

Addition of a binomial with a binomial :

Example 11 : Add 6x + 5 and 2x + 11.

Horizontal method	Vertical method
(6x + 5) + (2x + 11)	6x + 5
= 6x + 5 + 2x + 11	+ 2x + 11
= 6x + 2x + 5 + 11	8x + 16
(Taking like terms together)	
= (6+2)x + (5+11)	
= 8x + 16	


Example 12 : Add $3xy + 4y^2$ and $6xy - 5y^2$.

Vertical method

Example 13 : Add $9a^2 + 12$ and $-9 - 3a^2$.

Horizontal method	Vertical method
	$9a^2 + 12$
	$+ -3a^2 - 9$
	$6a^2 + 3$

MATHEMATICS 61 STD. 7




Horizontal method	Vertical method
$(6m^2 - 12) - (3m^2)$	$6m^2 - 12$
$= 6m^2 - 12 - 3m^2$	$-3m^2$
$= 6m^2 - 3m^2 - 12$	$3m^2 - 12$
$= (6 - 3) m^2 - 12$	
$= 3m^2 - 12$	

Example 15 : Subtract 4abc + 3 from 5abc - 7.

Horizontal method	Vertical method
(5abc - 7) - (4abc + 3)	
= 5abc - 7 - 4abc - 3	
= 5abc - 4abc - 7 - 3	
= (5 - 4) abc - 10	
= 1abc - 10	
= abc - 10	

MATHEMATICS 62 STD. 7



Addition of a trinomial with a monomial, a binomial and a trinomial :

Example 16 : Add $10x^2 - 5x + 2$ with $3x^2$.

Horizontal method	Vertical method
$(10x^2 - 5x + 2) + (3x^2)$	$10x^2 - 5x + 2$
$= 10x^2 - 5x + 2 + 3x^2$	$+3x^2+0+0$
$= 10x^2 + 3x^2 - 5x + 2$	$13x^2 - 5x + 2$
$= (10 + 3)x^2 - 5x + 2$	
$= 13x^2 - 5x + 2$	

Example 17 : Add $3m^2 - 2m + 7$ with 6m - 5.

MATHEMATICS

Horizontal method	Vertical method
$(3m^2 - 2m + 7) + (6m - 5)$	
$= 3m^2 - 2m + 7 + 6m - 5$	
$= 3m^2 - 2m + 6m + 7 - 5$	
$= 3m^2 + 4m + 2$	

Downloaded from https:// www.studiestoday.com

63

STD.7

and a contraction of the	Self to the Chickey of the self of the sel
6 : Polynomial	

Example 18 : Add $3a^2 + 20ab$	$+ b^2$ and $2a^2 - 6ab$ -	$-5b^2$.
---------------------------------------	----------------------------	-----------

Horizontal method	Vertical method
	$3a^2 + 20ab + b^2$
	$+ 2a^2 - 6ab - 5b^2$
	$5a^2 + 14ab - 4b^2$
Practice 6	

Add :

- (1) 2x + 3y + 5 and -7x (2) $12m^2 9m + 7$ and 3m 8
- (3) $2x^2 + 3x 5$ and $2x^2 4$ (4) 9b 10a + 15 and 3a + b + 2
- (5) 17a 13b 14 and 10a 9b 15 (6) $4p^2 3p 10$ and 30

• Subtraction of a monomial, a binomial and a trinomial with a trinomial :

Example 19 : Subtract 4m from $m^2 + 3m - 7$.

Horizontal method	Vertical method
$(m^2 + 3m - 7) - (4m)$	$m^2 + 3m - 7$
$= m^2 + 3m - 7 - 4m$	<u> </u>
$= m^2 + 3m - 4m - 7$	$m^2 - m - 7$
$= m^2 - 1m - 7$	
$= m^2 - m - 7$	

MATHEMATICS 64 STD. 7



Example 20 : Subtract $8x^2 - 4xy$ from $5x^2 - 4xy + y^2$.

Horizontal method	Vertical method
	$5x^2 - 4xy + y^2$
	$5x^{2} - 4xy + y^{2} + -8x^{2} + 4xy - 0y^{2}$
	$-3x^2 + 0xy + y^2$
	$= -3x^2 + y^2$


Example 21 : Subtract $7x^2 - 7x + 3$ from $10x^2 - 5x + 2$.

Horizontal method	Vertical method
$(10x^2 - 5x + 2) - (7x^2 - 7x + 3)$	
$= 10x^2 - 5x + 2 - 7x^2 + 7x - 3$	
$= 10x^2 - 7x^2 - 5x + 7x + 2 - 3$	
$= 3x^2 + 2x - 1$	



Subtract :

$5b^{2}$
x^2
_
$-2x^2$
4
27275



Downloaded from https:// www.studiestoday.com